

# DRAFT UGANDA STANDARD

First Edition  
2017

---

---

## Roasted Macadamia — Specification

---

---



Reference number  
DUS 1703: 2017

**Compliance with this standard does not, of itself confer immunity from legal obligations**

**A Uganda Standard does not purport to include all necessary provisions of a contract. Users are responsible for its correct application**

© UNBS 2017

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilised in any form or by any means, electronic or mechanical, including photocopying and microfilm, without prior written permission from UNBS.

Requests for permission to reproduce this document should be addressed to

The Executive Director  
Uganda National Bureau of Standards  
P.O. Box 6329  
Kampala  
Uganda  
Tel: +256 414 333 250/1/2/3  
Fax: +256 414 286 123  
E-mail: [info@unbs.go.ug](mailto:info@unbs.go.ug)  
Web: [www.unbs.go.ug](http://www.unbs.go.ug)

## Foreword

Uganda National Bureau of Standards (UNBS) is a parastatal under the Ministry of Trade, Industry and Cooperatives established under Cap 327, of the Laws of Uganda, as amended. UNBS is mandated to coordinate the elaboration of standards and is

- (a) a member of International Organisation for Standardisation (ISO) and
- (b) a contact point for the WHO/FAO Codex Alimentarius Commission on Food Standards, and
- (c) the National Enquiry Point on TBT Agreement of the World Trade Organisation (WTO).

The work of preparing Uganda Standards is carried out through Technical Committees. A Technical Committee is established to deliberate on standards in a given field or area and consists of key stakeholders including government, academia, consumer groups, private sector and other interested parties.

Draft Uganda Standards adopted by the Technical Committee are widely circulated to stakeholders and the general public for comments. The committee reviews the comments before recommending the draft standards for approval and declaration as Uganda Standards by the National Standards Council.

The committee responsible for this document is Technical Committee UNBS/TC 2, *[food and agriculture]*, Subcommittee SC 2, *[Edible oil seeds, fats and oils]*.

PUBLIC REVIEW DRAFT

# Roasted macadamia — Specification

## 1 Scope

This Draft Uganda Standard specifies the requirements, methods of sampling and testing for roasted macadamia of varieties (cultivars) grown from *Macadamia integrifolia*, *Macadamia tetraphylla* and *Macadamia ternifolia*, and their hybrids intended for direct consumption

## 2 Normative references

The following referenced documents referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

CODEX STAN 192, *General standard for food additives*

US EAS 35, *Edible salt — Specification*

EAS 38, *Labeling of prepackaged foods — Specification*

EAS 39, *Hygiene in the food and drink manufacturing industry — Code of practice*

US ISO 542, *Oilseeds -- Sampling*

US ISO 735, *Oilseed residues -- Determination of ash insoluble in hydrochloric acid*

US ISO 12193, *Animal and vegetable fats and oils -- Determination of lead by direct graphite furnace atomic absorption spectroscopy*

US ISO 15774, *Animal and vegetable fats and oils — Determination of cadmium content by direct graphite furnace atomic absorption spectrometry*

US ISO 17239, *Fruits, vegetables and derived products -- Determination of arsenic content -- Method using hydride generation atomic absorption spectrometry*

US ISO 7251, *Microbiology of food and animal feeding stuffs -- Horizontal method for the detection and enumeration of presumptive Escherichia coli -- Most probable number technique*

US ISO 6579, *Microbiology of food and animal feeding stuffs — Horizontal method for the detection of Salmonella spp.*

US ISO 16050, *Foodstuffs — Determination of aflatoxin B1, and the total content of aflatoxins B1, B2, G1 and G2 in cereals, nuts and derived products — High-performance liquid chromatographic method*

US ISO 27107, *Animal and vegetable fats and oils — Determination of peroxide value — Potentiometric end-point determination*

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

**3.1 roasted macadamia**  
 kernels which have been subjected to heat by dry or wet roasting and to which salt or other food ingredients may have been added

## 4 Requirements

### 4.1 General requirements

Roasted Macadamia shall have characteristic colour and be free from off-flavour

### 4.2 Raw materials

Raw Macadamia kernels for roasting shall conform to the standards for raw Macadamia kernels

### 4.3 Specific quality requirements

Roasted Macadamia shall comply with the requirements stipulated in Table 1.

**Table 1 — Limits for Roasted macadamia**

SN	Characteristic	limits	Test method
	Moisture content, % m/m, max.	1.5	US ISO 1026
	Salt as NaCl, % m/m, max.	2.0	US EAS 35
	Acid insoluble ash (roasted and salted), % m/m, max.	0.5	US ISO 735
	Peroxide value, max. , meq/kg	3	US ISO 3960

### 4.4 Grading

Roasted Macadamia may be graded. When graded the macadamia shall conform to the grades in Table 2.

**Table 2 — Grades for Roasted Macadamia**

S/N	Grade	Description	Whole kernel percentage	Size range (m/m)
i)	Premium	Whole kernels	100	25 to 15
ii)	Popular	Wholes and halves	min. 50	17 to 10
iii)	Cocktail	Wholes and halves	11 to 49	17 to 10
iv)	Medium	Halves and pieces	Max. 10	12 to 4
v)	Chips	Pieces and bits	None	5 to 0.5

<sup>a</sup>. Tolerance — Up to 5 % by mass of the next lower grade.

## 5 Food additives

Roasted Macadamia may be preserved by using food additives complying with CODEX STAN 192.

## 6 Contaminants

### 6.1 Pesticide residues

Roasted Macadamia shall conform to those maximum residue limits for pesticide residues established by Codex Alimentarius Commission for this commodity.

### 6.2 Heavy metals

Roasted macadamia shall be free from heavy metals in amounts which may represent a hazard to health. Heavy metal contaminants, if present, shall not exceed the limits stipulated in Table 3

**Table 3 — Heavy metal contaminant limits**

SL No.	Parameter	Maximum Limits	Test method
i)	Arsenic (As), ppm,	0.1	US ISO 17239
iii)	Lead (Pb), ppm,	0.10	US ISO 12193
iv)	Cadmium (Cd), ppm,	0.02	US ISO 15774

### 6.3 Aflatoxin

Roasted macadamia kernels shall conform to those maximum limits for aflatoxin stated in Table 4

**Table 4- Limits for aflatoxin in roasted macadamia kernels**

SN	Aflatoxin type	Limits, $\mu\text{g}/\text{kg}$ , max.	Methods of test
1	Aflatoxin B <sub>1</sub>	5	US ISO 16050
2	Total Aflatoxins	10	

## 7 Hygiene

Roasted Macadamia shall be handled in hygienic manner in accordance with EAS 39. Roasted Macadamia shall conform to the microbiological limits in Table 5

**Table 5 — Microbiological limits for Roasted Macadamia**

SL No.	Micro-organism	Maximum limits	Test method
i)	<i>Escherichia coli</i> , cfu/g	Absent	US ISO 7251
ii)	<i>Salmonellae</i> , in 25g	Absent	US ISO 6579

## 9 Packaging

9.1 Roasted Macadamia may be sold packaged or loose. Packaged macadamia kernels shall be packaged in such a way to protect the produce from mechanical, heat and frost damage. Macadamia kernels shall be packaged in food grade materials that will safeguard the hygienic, nutritional, technological and organoleptic qualities of the produce.

9.2 The packaging materials shall conform to the environmental legislation of the destination country.

## 10 Weights and measures

The weight of the package of the product shall comply with Weights and Measures regulations of the importing Partner State.

## 11 Labelling

11.1 In addition to the requirements of EAS 38, the following specific requirements shall apply and shall be legibly and indelibly marked:

- a) name of the product shall be "Roasted Macadamia
- b) name of the product shall be "Roasted Macadamia";
- c) name and location address of the exporter and/or packer shall be declared;
- d) country of origin;
- e) Net weight;
- f) lot identification (batch number);
- g) declaration of ingredients, if used;
- h) expiry date;
- i) grade; and
- j) storage instructions

## 12 Sampling

Sampling shall be done in accordance to US ISO 542.



## Annex A (normative)

### Annex title

#### A.1 General

##### Procedure

**A.1.1** Weigh accurately about 10 g of the material in a suitable moisture dish previously dried in an electric oven and weighed. Place the dish in a vacuum oven at 95 °C to 100 °C under pressure 100 mm/Hg. Cool the dish in a desiccator and weigh with the lid on. Repeat the process of heating, cooling and weighing at half- hour intervals until the loss in weight between two successive weighings is less than 1 mg. Record the lowest weight obtained.

**A.1.2** Calculation and expression of results, moisture % by mass

$$= \frac{(m_1 - m_2) \times 100}{m_1 - m_3}$$

where:

$m_1$ : is the mass in g of the dish and sample before drying;

$m_2$ : is the mass in g of the dish and sample after drying; and

$m_3$ : is the mass in g of the dish only

## Annex B (normative)

### Determination of oil content

#### B.1 Reagents

B.1.1 Petroleum ether of boiling range of 40 °C to 60 °C.

B.1.2 Hexane of food grade

#### B.2 Procedure

Weigh accurately about 2.5 g of the sample, dried as described under A1.1, and extract with petroleum ether or hexane, food grade, in a Soxhlet or other suitable extractor. The extraction period may vary from 4h at a condensation rate of 5.6 drops per second. Dry the extract on a steam bath for 30 min, cool in a desiccator and weigh. Continue at 30 min intervals this alternative drying and weighing until the difference between two successive weighings is less than 1 mg.

Note the lowest mass.

#### B.3 Calculation and expression of results

Crude fat (on moisture-free basis), % by mass

$$= 100 \frac{(m_1 - m_2)}{m_3}$$

where,

$m_1$ : is the mass in g of the extraction flask with dried extract;

$m_2$ : is the mass in g of extraction flask; and

$m_3$ : is the mass in g of the dried sample for the test

## Annex C (normative)

### Determination of free fatty acid

#### C.1 Apparatus

Soxhlet fat extraction apparatus.

#### C.2 Reagents

**C.2.1** Petroleum, ether, distilling below 65 °C, or ethyl ether.

**C.2.2** Alcohol potassium hydroxide, 0.1N (use absolute or alcohol denatured with Me(OH)).

**C.2.3** Alcohol-ether mixture, equal volumes of 96 % alcohol and ethyl ether.

**C.2.4** Phenolphthalein solution, one % in alcohol ether and add alcoholic KOH solution to a faint pink.

#### C.3 Procedure

**C.3.1** Extract 10.00 ± 0.01 g of the sample taken in a thimble with petroleum ether for about 16 h in a Soxhlet extraction apparatus. Completely evaporate the solvent from extraction flask (weighed previously) on a steam bath, cool and weigh the Soxhlet flask with the residue and from the difference of the mass of fat obtained from the material. Dissolve the residue in the extraction flask with 50 ml of the alcohol-ether phenolphthalein solution. Titrate the dissolved extract, with standard potassium hydroxide solution to a faint pink colour which persists for 10s. If emulsion is formed during titration, dispel by adding second 50 ml portion of the alcohol-ether phenolphthalein solution.

**C.3.2** Make a blank titration on 50 ml of the alcohol-ether phenolphthalein solution and subtract this value from the titration value of the sample. If the additional 50 ml portion of the alcohol-ether phenolphthalein solution is added, double the blank titration.

## Annex D (normative)

### Determination of salt

#### D.1 Reagents

- D.1.1 Acetone.
- D.1.2 10 % calcium acetate solution.
- D.1.3 HNO<sub>3</sub>.
- D.1.4 0,1 N AgNO<sub>3</sub>.

D.1.5 FeAlum indicator.

D.1.6 0.1 N NH<sub>4</sub>SCN.

#### D.2 Procedure

D.2.1 Weight 2 g of thoroughly mixed sample into a platinum or silica dish.

D.2.2 Disperse sample with 10 ml of acetone.

D.2.3 Remove acetone at room temperature, with air current.

D.2.4 Add, and thoroughly mix, 10 ml of 10 % calcium acetate.

D.2.5 Carefully dry on a steam bath.

D.2.6 Ash in a muffle oven at 550 °C (1022°F) (complete ashing is not necessary).

D.2.7 Place ash in a beaker and dissolve ash in 25 ml HNO<sub>3</sub> (1 + 3).

D.2.8 Add at least 2 - 4 ml of 0.1 N AgNO<sub>3</sub> that is just enough to precipitate all chloride present.

D.2.9 Add at least 5 ml of 0.1 N AgNO<sub>3</sub> in excess, to 8.

D.2.10 Heat to boil, cool, then add 5 ml FeAlum indicator.

D.2.11 Titrate excess Ag with 0.1 N NH<sub>4</sub>SCN (which has been standardized to equalize normalities) to a permanent light brown end point

D.2.12 Subtract the amount of NH<sub>4</sub> SCN used (D.2.11) from the total AgNO<sub>3</sub> used in D.2.8 and D.2.9. The resulting difference is the ml of 0.1 N AgNO<sub>3</sub> used in the calculation of salt (D.2.13).

D.2.13 Calculate as % NaCl

$$\frac{(\text{mL of 0.1 N AgNO}_3 - 0.05845)(100)}{100}$$

## Bibliography

- [1] RS 171, *Roasted macadamia — Specification*

PUBLIC REVIEW DRAFT

## Certification marking

Products that conform to Uganda standards may be marked with Uganda National Bureau of Standards (UNBS) Certification Mark shown in the figure below.

The use of the UNBS Certification Mark is governed by the Standards Act, and the Regulations made thereunder. This mark can be used only by those licensed under the certification mark scheme operated by the Uganda National Bureau of Standards and in conjunction with the relevant Uganda Standard. The presence of this mark on a product or in relation to a product is an assurance that the goods comply with the requirements of that standard under a system of supervision, control and testing in accordance with the certification mark scheme of the Uganda National Bureau of Standards. UNBS marked products are continually checked by UNBS for conformity to that standard.

Further particulars of the terms and conditions of licensing may be obtained from the Director, Uganda National Bureau of Standards.





PUBLIC REVIEW DRAFT

---

---

ICS nn.nnn.nn

Price based on nn pages