



## **DRAFT EAST AFRICAN STANDARD**

---

**Sodium hypochlorite solutions for domestic and industrial use —  
Specification**

**EAST AFRICAN COMMUNITY**

---

### Copyright notice

This EAC document is copyright-protected by EAC. While the reproduction of this document by participants in the EAC standards development process is permitted without prior permission from EAC, neither this document nor any extract from it may be reproduced, stored or transmitted in any form for any other purpose without prior written permission from EAC.

Requests for permission to reproduce this document for the purpose of selling it should be addressed as shown below or to EAC's member body in the country of the requester:

© East African Community 2020 — All rights reserved  
East African Community  
P.O. Box 1096,  
Arusha  
Tanzania  
Tel: + 255 27 2162100  
Fax: + 255 27 2162190  
E-mail: [eac@eachq.org](mailto:eac@eachq.org)  
Web: [www.eac-quality.net](http://www.eac-quality.net)

Reproduction for sales purposes may be subject to royalty payments or a licensing agreement. Violators may be prosecuted.

## Foreword

Development of the East African Standards has been necessitated by the need for harmonizing requirements governing quality of products and services in the East African Community. It is envisaged that through harmonized standardization, trade barriers that are encountered when goods and services are exchanged within the Community will be removed.

The Community has established an East African Standards Committee (EASC) mandated to develop and issue East African Standards (EAS) and other deliverables. The Committee is composed of representatives of the National Standards Bodies in Partner States, together with the representatives from the public and private sector organizations in the community.

East African Standards are developed through Technical Committees that are representative of key stakeholders including government, academia, consumer groups, private sector and other interested parties. Draft East African Standards are circulated to stakeholders through the National Standards Bodies in the Partner States. The comments received are discussed and incorporated before finalization of standards, in accordance with the Principles and procedures for development of East African Standards.

East African Standards and other deliverables are subject to review, to keep pace with technological advances. Users of the East African Standards are therefore expected to ensure that they always have the latest versions of the standards they are implementing.

The committee responsible for this document is Technical Committee EASC/TC 074, *Surface active agents*

Attention is drawn to the possibility that some of the elements of this document may be subject of patent rights. EAC shall not be held responsible for identifying any or all such patent rights.

This second edition cancels and replaces the first edition (EAS 295:2002), which has been technically revised.



## Sodium hypochlorite solutions for domestic use — Specification

### 1. Scope

This Draft East African Standard specifies requirements, sampling and test methods for sodium hypochlorite intended for domestic and Industrial use.

### 2. Normative references

Not applicable

### 3. Terms and definitions

For the purpose of this draft standard the following definitions apply.

#### 3.1

##### available chlorine

quantity of chlorine chemically equivalent to the oxygen that would be released during the complete decomposition of the, sodium hypochlorite to sodium chloride and oxygen

NOTE - This is a conventional way of expressing the concentration of sodium hypochlorite solution. The available chlorine is 0.95 times the sodium hypochlorite content and is a measure of the oxidising power of sodium hypochlorite solutions.

#### 3.2

##### nominal concentration

minimum available chlorine content of the sodium hypochlorite solution under test, at the time of manufacture

#### 3.3

##### product unit

unit of the final product, packed in a suitable container

#### 3.4

##### lot

number of containers consisting of product of the same type and style, which have been manufactured and packed under essentially the same conditions

### 4. Requirements

#### 4.1. General requirements

4.1.1 Sodium hypochlorite solutions shall be of one of the following nominal concentrations, as required

- a) 15.0 % m/v for industrial use;
- b) 5.0 % m/v for domestic use; or
- c) 3.5 % m/v for domestic use.

4.1.2 Sodium hypochlorite solution shall be a clear liquid, free from sediment and suspended matter. A solution shall be considered to be clear if the small amount of salts that has crystallised from the solution dissolves completely when the solution is mixed with twice its volume of distilled water. When so, required, the solution shall contain laundry blue. The laundry blue may settle on standing but shall disperse completely in the solution when the solution, in the original container, is shaken for 30 s.

#### 4.2. Specific requirements

The solution shall comply with the specific requirements given below in Table 1 for domestic use and Table 2 for industrial use

Table 1 — **Specific requirements for a sodium hypochlorite for domestic use**

S/No:	Characteristic	Requirement	Test method
i	Available chlorine content determined within 14 days $\pm$ 2 d of the date of manufacture, % (m/v), min 5.0 % nominal concentration 3.5 % nominal concentration	5.0 3.5	Annex A
ii	Available chlorine content determined from 14 <sup>th</sup> to the 60 <sup>th</sup> day $\pm$ 2 d of the date of manufacture, % (m/v), min 5 % nominal concentration 3.5 % nominal concentration	4.5 3.2	Annex A
iii	Sodium hydroxide content, % (m/v), max	0.5	Annex B
iv	Sediment content, % (mass fraction expressed as a percentage), max	0.1	Annex C

Table 2 - **Specific requirements for sodium hypochlorite for Industrial use**

S/No:	Parameter	Requirement	Test method
i	Available chlorine content determined on the date of manufacture, % (m/v), min	15	Annex A
ii	Available chlorine content determined on the 14 <sup>th</sup> day $\pm$ 2 d after date of manufacture, % (m/v), min	13	Annex A
iii	Sodium hydroxide content, % (m/v), max	1.5	Annex B
iv	Sediment content, % (mass fraction expressed as a percentage), max	0.1	Annex C

### 5. Packing and Labelling

#### 5.1. Packing

Sodium hypochlorite shall be packed in a suitable opaque containers that withstand normal handling and transportation and that will prevent leakage and contamination of the product.

#### 5.2. Labelling

Each container shall be in prominent, legibly and indelibly labelled either in English, Kiswahili or French or combination or any other language as agreed between the manufacturer and supplier with the following information:

- a) the name of product as "Sodium hypochlorite for domestic use" or "Sodium hypochlorite for industrial use";
- b) nominal available chlorine content and/or the percentage composition of the solution;
- c) name and physical address of the manufacturer and trade mark if any;
- d) net contents;
- e) the words "**store in a cool place away from direct sunlight; avoid contact With aluminium, zinc, tin and their alloys and do not mix with acids**";
- f) the instructions for use;
- g) country of origin;
- h) date of manufacture and best before date;

i) Caution:

- "Keep out of reach of children", avoid contact with eyes, or words having similar meaning;
- Do not use on wool, silk, rayon, and leather";

NOTE - Sodium hypochlorite solution may have deleterious effect on certain resin-treated materials, such as grease resistant, dry drip, embossed and glazed fabrics. Coloured fabrics may lose their colour if their dyes are not colour fast to hypochlorite

## 6. Sampling and compliance with the standard

### 6.1. Sampling

6.1.1. For ascertaining the conformity of the lot to the requirements of this standard, tests shall be carried out on each lot separately.

6.1.2. The number of packages and product units from each container respectively to be selected for drawing the sample shall be in accordance with 1.

6.1.3. The sample so drawn shall be deemed to represent the lot. From a given lot, for product units of 500 cm<sup>3</sup> or more a sample of one unit shall be chosen.

6.1.4. After checking the lot for compliance with the relevant requirements of 5.1 and 5.2, take from it at random the number of containers shown in column 2 of Table 1, relative to the appropriate lot size given in column 1

6.1.5. Reserve half the containers for the determination of the characteristics other than stability and the other for the determination of stability.

**Table 1 — Scale of sampling**

1	2	3
Lot size, number of cartons (packages) in a lot	Number of cartons (containers) to be selected	Number of product units to be selected from each carton
25 – 50	4	4
51 – 100	6	6
101 – 500	8	2
501 – 1500	10	1
1501 – 5000	12	1

6.1.6. The packages (cartons) shall be selected at random, using tables of random numbers. If these are not available; the following procedure shall be applied:

Starting from any package, count all the packages in one order as 1, 2, 3... N, selecting every k<sup>th</sup> package, where k is the integral part of  $N \div n$ .

From each package thus selected, draw at random an equal number of cakes so as to obtain a total mass of at least 2 kg.

### 6.2. Compliance with the standard

The lot shall be deemed to comply with the requirements of the standard if after inspection of the containers and testing of the sample taken in accordance with 5.1 no defective is found.





## Annex A (Normative)

### Determination of available chlorine

#### A.1 Method 1

##### A.1.1 Principle

Known volumes of sodium hypochlorite solution and arsenite solution are mixed and titrated against iodine solution.

##### A.1.2 Reagents

###### A.1.2.1 Phenolphthalein indicator solution,

Dissolve 1 g of phenolphthalein in 100 mL of ethanol (95 % v/v)

###### A.1.2.2 Starch solution

Prepare the starch solution immediately before use as follows

Make a paste of 1 g of soluble starch and a small amount of distilled water. Pour the paste into 100 ml of boiling water, stirring it constantly. Boil for approximately 1 min and cool.

###### A.1.2.3 Standard sodium arsenite solution (c (NaAsO<sub>2</sub>) = 0.025 mol/l)

Dissolve exactly 4.946 g of arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) in 150 mL of warm water containing about 3 g of sodium hydroxide. Cool, add one drop of phenolphthalein indicator, and then neutralise the excess alkali with dilute sulphuric acid (1:3) until the solution is just decolourised. Transfer qualitatively to a 1 litre volumetric flask and add 500 ml of water containing about 25 g of sodium bicarbonate. If a pink colour develops, add dilute sulphuric acid drop by drop until the solution becomes colourless. Make up to the mark with water, and mix well.

###### A.1.2.4 Standard iodine solution 0.025M.

###### A.1.2.4.1 Preparation:

Mix 6.35 g ± 0.05 g of Iodine with twice this mass of potassium iodide. Dissolve the mixture in 100 mL of water and then filter the solution through a filter paper of medium porosity, into a 1 L volumetric flask. Wash the filter paper three times with 25 mL portions of water. Make up to the mark with water and mix well.

###### A.1.2.4.2 Standardisation

Standardise the iodine solution as follows:

- Pipette 20 mL of the standard sodium arsenite solution (A.1.2.3) into a 250 mL conical flask;
- Add 5 mL of water and about 5 g of bicarbonate and shake well the flask until the sodium bicarbonate has completely dissolved;
- While shaking the flask continuously, titrate the solution with the iodine solution until the yellow
- colour formed disappears slowly from the solution when it is shaken; and
- Add 5 mL of the starch solution (A.1.2.2) and continue the titration until a permanent blue colour is obtained.

Calculate the concentration of the standard iodine solution  $M$ , in moles per litre, as follows:

$$M = 1.0/V$$

Where

$V$  is the volume of iodine solution used in the titration, in millimetres.

### A.1.3 Procedure

**A.1.3.1** Pipette 25 mL of sodium hypochlorite solution into a volumetric flask containing about 500 mL of distilled water, dilute to 1000 mL mark and mix well

**A.1.3.2** Pipette 25 mL this solution and 25 mL of the standard sodium arsenite solution into a 250 mL conical flask.

**A.1.3.3** Add 5 g of sodium bicarbonate and shake the flask until it has completely dissolved.

**A.1.3.4** Titrate with the standard iodine solution until the yellow colour formed slowly disappears from the solution when it is shaken.

**A.1.3.5** Add 5 ml of the starch solution and continue the titration until a permanent blue colour is obtained.

### A.1.4 Calculation

Calculate available chlorine content  $C$ , in grams per litre, as follows

$$C = ((1.25 - VM) \times 2836.0) V_1$$

where,

$V$  is the volume of iodine solution used in the titration in millilitres;

$V_1$  is the volume of sodium hypochlorite (test specimen taken) in millilitres; and

$M$  is the concentration of the standard iodine solution, in moles per litre.

## A.2 Method 2 (Alternative method)

### A.2.1 Principle

The sample is added to acidified solution of potassium iodide and the released iodine is titrated with standard sodium thiosulphate solution to the usual starch end point.

### A.2.2 Reagents

**A.2.2.1** Acetic acid, glacial

**A.2.2.2** Potassium iodide (KI), crystals, iodate free

**A.2.2.3** Sodium thiosulphate ( $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ ), standard solution - Dissolve 25 g of  $\text{Na}_2\text{SO}_3$  crystals in freshly boiled and cooled water and dilute to 1 L. The solution is more stable if the glassware is cleaned with sulphuric-chromic acid and thoroughly rinsed with water. Standardize against potassium iodate ( $\text{KIO}_3$ ) as follows: Weigh out accurately 3.567 g of dry  $\text{KIO}_3$  and transfer to a 1 L volumetric flask. Dissolve with water, make up to the mark and mix thoroughly. This solution will be exactly 0.1000 N. To standardize the  $\text{Na}_2\text{S}_2\text{O}_3$  solution, carefully pipette a 50 mL aliquot of the  $\text{KIO}_3$  solution into a 250 mL Erlenmeyer flask and dilute to 100 mL with water. Add 1 g of KI crystals. When it is dissolved, add 15 mL of 1.0 N hydrochloric acid and titrate immediately with the  $\text{Na}_2\text{S}_2\text{O}_3$  solution. When the solution becomes light yellow, add 1 mL of starch indicator solution and complete the titration to the disappearance of the blue colour. Standardize at least monthly.

Calculate the normality of the  $\text{Na}_2\text{S}_2\text{O}_3$  solution as follows:

$$\text{Normality, } N1 = (50 \times 0.1)/A$$

where,

$A$  is the value of  $\text{Na}_2\text{S}_2\text{O}_3$  solution required for titration of  $\text{KIO}_3$  solution.

**A.2.2.4** Starch indicator solution, mix 0.5 g of soluble starch with 5 mL of cold water and add to 95 mL of boiling water. Mix, cool and store in a sterilized bottle. Replace frequently or add 0.1 % salicylic acid to minimize deterioration.

**A.2.3 Procedure**

Dissolve 2 g to 3 g of KI crystals to 50 mL of water in a 250 mL Erlenmeyer flask. Add 10 mL of acetic acid. Then pipette the aliquot of sample into the solution keeping the tip of the pipette beneath the surface of the solution until drained. Titrate at once with 0.1 N Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution until the iodine colour is nearly gone then add 1 mL of starch indicator solution and complete the titration to the disappearance of the iodine colour. Record the titration as A.

**A.2.4 Calculations**

**A.2.4.1** Calculate the available chlorine as follows:

$$\text{Available chlorine as Cl, g/L} = \frac{A \times N_1 \times 35.46}{V}$$

**A.2.4.2** Calculate the sodium hypochlorite content as follows;

$$\text{Sodium hypochlorite (NaOCl), g/L} = \frac{A \times N_1 \times 37.22}{V}$$

where

A is the volume, in millilitres, of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution required for titration of the sample;  
N<sub>1</sub> is the normality of the Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution; and  
V is the volume, in millilitres, of original sample in aliquot used.

## Annex B (Normative)

### Determination of sodium hydroxide

#### B.1 Principle

The sodium hypochlorite is decomposed with hydrogen peroxide, carbonate precipitated with barium chloride and the hydroxide determined by titration with hydrochloric acid.

#### B.2 Reagents

**B.2.1 Barium chloride solution** 100 g of barium chloride ( $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ ) in water and dilute to 1 L with water. Filter the solution of turbid.

**B.2.2 Hydrogen peroxide solution** - approximately 3 % (10 volumes)

**B.2.3 Sodium hydroxide solution** - approximately 4 g/L in water

**B.2.4 Phenolphthalein indicator** - prepared as described in 6.1.2.1

**B.2.5 Screened methyl orange indicator** - Dissolve 0.2 g of methyl orange and 0.28 g of xylene cyanol FF in 100 mL of ethanol (50 %).

**B.2.6 Standard hydrochloric acid solution** 0.1M

#### B.3 Procedure

**B.3.1** Place 50 ml of barium chloride solution and 40 ml of hydrogen peroxide solution in a 250 ml conical flask.

**B.3.2** Add 10 drops of phenolphthalein indicator and then add sodium hydroxide solution drop by drop until a permanent faint pink colour is obtained.

**B.3.3** Immediately pipette 10 ml of sodium hypochlorite solution drop by drop into the flask, taking care that the effervescence does not become excessive.

**B.3.4** When the effervescence subsides, shake the flask vigorously for 1 min.

**B.3.5** Add another drop of phenolphthalein indicator and rapidly titrate the, solution with the standard hydrochloric acid solution (B.2.6) until the pink colour first disappears. Do not continue the titration if the pink colour reappears on standing.

#### B.4 Calculation

Calculate the sodium hydroxide content  $X$ , in grams per litre, as follows:

$$X = V \times M \times 4.0$$

Where

$V$  is the volume, of standard hydrochloric acid solution used in the titration in millilitres

$M$  is the concentration of the standard hydrochloric acid solution, in moles per litre.

**Annex C**  
(Normative)

**Sediment content**

**C.1 Procedure**

**C.1.1** Mix the test specimen of sodium hypochlorite solution thoroughly and then accurately weigh out approximately 300 g into a 500 mL beaker.

**C.1.2** Filter the solution through a Whatman GF/A glass-fibre filter paper that has been dried at 100 °C ± 2 °C and tared, or equivalent.

**C.1.3** Wash the beaker and the sediment five times with 20 mL portions of cold water and then dry the glass fibre filter paper with the sediment at 100 °C ± 2 °C until a constant mass is attained.

**C.2 Calculation**

Calculate the sediment content, *S*, as a percentage, as follows:

$$S = \frac{m_1}{m_2} \times 100$$

where

*m*<sub>2</sub> is the mass of the sediment after it has been dried, in grams;

*m*<sub>1</sub> is the mass of test specimen taken, in grams.