

Fluoxastrobin

Commodity	MRL (draft) ppm	MRL (current) ppm
Wheat	○ 0.2	
Barley	○ 0.4	
Corn (maize, including pop corn and sweet corn)	○ 0.02	
Other cereal grains ¹	○ 2	
Soybeans, dry	○ 0.05	
Beans, dry ²	○ 0.2	
Peas	○ 0.2	
Broad beans	○ 0.2	
Other pulses ³	○ 0.2	
Potato	0.01	0.01
Apple	○ 1	
Japanese pear	○ 0.6	
Pear	○ 0.6	
Strawberry	2	2
Grape	○ 2	
Rapeseeds	○ 0.7	
Cattle, muscle	○ 0.05	
Pig, muscle	○ 0.02	
Cattle, fat	○ 0.1	
Pig, fat	○ 0.03	
Cattle, liver	○ 0.2	
Pig, liver	○ 0.06	
Cattle, kidney	○ 0.2	
Pig, kidney	○ 0.06	
Cattle, edible offal ⁴	○ 0.2	
Pig, edible offal	○ 0.06	
Milk	○ 0.03	

○ : Commodities for which MRLs are to be raised.

Note: The residue definition for agricultural products is sum of Fluoxastrobin and metabolite Z-isomer [(Z)-{2-[6-(2-chlorophenoxy)-5-fluoropyrimidin-4-yl]phenyl}(5,6-dihydro-1,4,2-dioxazin-3-yl)methanone O-methyl oxime], expressed as Fluoxastrobin. The residue definition for animal products is sum of Fluoxastrobin, metabolite Z-isomer, and metabolite M55[6-(2-chlorophenoxy)-5-fluoro-4-pyrimidol], expressed as Fluoxastrobin.

* The uniform limit 0.01 ppm will be applied to commodities not listed above.

1. "Other cereal grains" refers to all cereal grains, except rice (brown rice), wheat, barley, rye, corn (maize) and buckwheat.

2. "Beans, dry" includes butter beans, cowbeans (red beans), lentil, kidney beans, lima beans, pedia, sultani, sultapya and white beans.
3. "Other pulses" refers to all legumes/pulses, except soybeans (dry), beans (dry), peas, broad beans, peanuts (dry) and spices.
4. "Edible offal" refers to all edible parts, except muscle, fat, liver and kidney.